

# The Wag is back.

## Benefits for you and your dog.

- ✓ Helps control pain and inflammation associated with
  - Osteoarthritis
  - After orthopedic or dental surgery
- ✓ Affordable pain control
- ✓ Tasty, beef-flavored chewable tablets
- ✓ For all dogs of all sizes

Doxidyl™ (deracoxib) is a highly effective NSAID (non-steroidal, anti-inflammatory drug) that has been trusted by veterinarians for over 15 years.

## Doxidyl™ (deracoxib) Chewable Tablets

- ✓ Inhibit the production of the enzyme that contributes to inflammation
- ✓ Best given with food, but tasty enough to be given without
- ✓ Scored to allow for precise dosing
- ✓ Come in 4 dosing sizes: 12 mg, 25 mg, 75 mg, and 100 mg



See your dog wagging again with  
**Doxidyl™ (deracoxib)  
Chewable Tablets,**  
tasty pain control for osteoarthritis  
at an affordable price.



Your dog doesn't have to suffer from osteoarthritis pain and inflammation. Tasty, affordable, and effective pain control in a chewable tablet.

Doxidyl (deracoxib) will have your dog wagging again.

**Ask your veterinarian about Doxidyl™ (deracoxib) today.**



**Learn more at [Doxidyl.com](http://Doxidyl.com)**

#### Important Safety Information:

For use in dogs only. Do not use in cats. As a class, nonsteroidal anti-inflammatory drugs (NSAIDs) may be associated with gastrointestinal, kidney or liver side effects. These are usually mild but may be serious. If side effects occur, pet owners should discontinue use and contact their veterinarian. As with any NSAID, all dogs should undergo a thorough history and physical examination before the initiation of NSAID therapy. Concurrent use with other anti-inflammatory drugs, such as NSAIDs or corticosteroids, should be avoided. The safe use of DOXIDYL™ (deracoxib) Chewable Tablets in dogs younger than 4 months of age, dogs used for breeding or in pregnant or lactating dogs has not been evaluated. Refer to the full prescribing information for complete details.



**Chewable Tablets**

**For Oral Use in  
Dogs Only  
Do Not Use in Cats**

#### Information for Dog Owners

DOXIDYL Chewable Tablets are for the control of pain and inflammation due to osteoarthritis or following orthopedic and dental surgery.

This summary contains important information about DOXIDYL tablets. You should read this information before starting your dog on DOXIDYL tablets. This sheet is provided only as a summary and does not take the place of instructions from your veterinarian. Talk to your veterinarian if you do not understand any of this information or you want to know more about DOXIDYL tablets.

##### What is DOXIDYL?

DOXIDYL tablets are a prescription non-steroidal anti-inflammatory drug (NSAID) of the coxib class. They are indicated for the control of postoperative pain and inflammation associated with orthopedic (bone) and dental surgery in dogs and for the control of pain and inflammation (soreness) associated with osteoarthritis in dogs. The tablets are flavored to make administration more convenient.

##### What kind of results can I expect when my dog takes DOXIDYL tablets for postoperative orthopedic and dental pain and inflammation?

DOXIDYL tablets allow your dog to recover more comfortably by controlling pain and inflammation that follow orthopedic and dental surgery.

- The postoperative orthopedic pain dose is a higher dose; administration at this dose **should not exceed 7 days total, including the days the patient is administered DOXIDYL while in the hospital.**

- Control of pain and inflammation may vary from dog to dog.
- If DOXIDYL tablets are not given according to your veterinarian's directions, your dog's pain may return.

- Consult your veterinarian if your dog appears to be uncomfortable.

##### What kind of results can I expect when my dog takes DOXIDYL tablets for pain and inflammation due to osteoarthritis?

Osteoarthritis is a painful condition caused by damage to cartilage and other parts of the joint that may result in the following changes or signs in your dog:

- Limping or lameness
- Decreased activity or exercise (reluctance to stand, climb stairs, jump or run, or difficulty in performing these activities)
- Stiffness or decreased movement of joints

While DOXIDYL is not a cure for osteoarthritis, it can control the pain and inflammation of osteoarthritis and improve your dog's mobility. Response may vary from dog to dog but can be quite dramatic. **DOXIDYL tablets may need to be given on a periodic basis for the animal's lifetime. Use the lowest dose to provide adequate relief. Always consult with your veterinarian before altering the dose.**

##### What dogs should not take DOXIDYL tablets?

Your dog should not be given DOXIDYL tablets if s/he:

- Has had an allergic reaction to deracoxib, the active ingredient in DOXIDYL tablets
- Has had an allergic reaction (such as hives, facial swelling, or red or itchy skin) to aspirin or other NSAIDs
- Is presently taking aspirin, other NSAIDs, or corticosteroids (unless directed by your veterinarian)
- Has bloody stool or vomit
- Has a pre-existing kidney or liver condition
- Has any condition predisposing to dehydration
- Is anorexic (loss of appetite)

##### DOXIDYL tablets should only be given to dogs. Do not use in cats.

People should not take DOXIDYL tablets. Keep DOXIDYL tablets and all medication out of reach of children. Call your physician immediately if you accidentally take DOXIDYL tablets.

##### What to discuss with your veterinarian before giving DOXIDYL tablets?

Tell your veterinarian about:

- Any side effects your dog has experienced from DOXIDYL tablets or other NSAIDs
- Any digestive upset (vomiting or diarrhea) your dog has had
- Any kidney disease your dog has had
- Any other medical problems or allergies that your dog has now or has had in the past
- All medications that you are giving your dog or plan to give your dog, including those you can get without prescription and any dietary supplements

- If you plan to breed your dog, or if your dog is pregnant or nursing  
Talk to your veterinarian about:

- The orthopedic or dental surgery your dog will undergo
- What tests might be done before surgery is performed or DOXIDYL tablets are prescribed
- The signs of pain or inflammation that may occur after surgery
- Normal events that can be expected after your dog undergoes surgery
- The proper amount of exercise after surgery to aid recovery
- The signs of osteoarthritis you have observed (for example, limping or stiffness)
- The importance of weight control, physical therapy and exercise in the management of osteoarthritis
- How often your dog may need to be examined by your veterinarian
- The risks and benefits of using DOXIDYL tablets

##### How to give DOXIDYL tablets to your dog.

DOXIDYL tablets should be given according to your veterinarian's instructions. Your veterinarian will tell you what amount of DOXIDYL tablets is right for your dog and for how long they should be given. Do not change the way you give DOXIDYL tablets to your dog without first speaking with your veterinarian. DOXIDYL tablets should be given by mouth and may be given with or without food, although with food is preferable.

##### What are the possible side effects that may occur in my dog during therapy with DOXIDYL tablets?

DOXIDYL tablets may cause some side effects in individual dogs. Serious side effects associated with this drug can occur with or without warning and, in some cases, result in death. The most common side effects associated with DOXIDYL therapy involve the digestive tract (vomiting, decreased appetite and diarrhea). Liver and kidney problems have also been reported. It is important to stop the medication and contact your veterinarian immediately if you think your dog may have a medical problem or side effect while on DOXIDYL tablets. If you have additional questions about possible side effects, talk with your veterinarian or call Ceva Animal Health, LLC at 1-800-999-0297.

Look for the following side effects that may indicate that your dog is having a problem with DOXIDYL tablets or may have another medical problem:

- Vomiting
- Decrease in appetite
- Change in behavior, such as depression, restlessness, aggression or apprehension
- Change in bowel movements such as diarrhea or change in stool color (black, tarry or bloody stool)
- Change in drinking or urination
- Yellowing of gums, skin or whites of the eyes (jaundice)

##### Can DOXIDYL tablets be given with other medications?

DOXIDYL tablets should not be given with other non-steroidal anti-inflammatory drugs (NSAIDs) or corticosteroids (for example, aspirin, carprofen, etodolac, prednisone).

Tell your veterinarian about all medications that you have given your dog in the past, and any medications that you are planning to give with DOXIDYL tablets. This should include any medications that you can get without a prescription and any dietary supplements. Your veterinarian may want to evaluate the potential for any drug interactions and to assure drug compatibility.

##### What can I do in case my dog eats more than the prescribed amount of DOXIDYL tablets?

Contact your veterinarian immediately if your dog eats more than the prescribed amount of DOXIDYL tablets.

##### What else should I know about DOXIDYL tablets?

This sheet provides a summary of information about DOXIDYL tablets. If you have any questions or concerns about DOXIDYL tablets, postoperative orthopedic and dental pain and inflammation, or osteoarthritis, talk to your veterinarian.

As with all prescribed medications, DOXIDYL tablets should only be given to the dog for which they are prescribed. They should be given to your dog only for the condition for which they were prescribed, at the prescribed dose, as directed by your veterinarian. It is important to periodically discuss your dog's response to DOXIDYL tablets at regular checkups. Your veterinarian will best determine if your dog is responding as expected and if your dog should continue receiving DOXIDYL tablets.

Approved by FDA under ANADA # 200-637  
©2019 Ceva Animal Health, LLC



Doxidyl™ trademark is the property of Ceva Santé Animale S.A.