

### **Meloxidyl® (meloxicam) Injection ISI for dogs:**

As a class, cyclo-oxygenase inhibitory NSAIDs may be associated with gastrointestinal, kidney, or liver side effects. The most common side effects reported in field studies were vomiting and soft stool/diarrhea. These are usually mild but may be serious. If side effects occur, pet owners should contact their veterinarian. Pets should be evaluated for pre-existing conditions and currently prescribed medications prior to treatment with MELOXIDYL, then monitored regularly while on treatment. Concurrent use with another NSAID, corticosteroid, or nephrotoxic medication should be avoided. The safe use of MELOXIDYL in dogs younger than 6 months of age, dogs used for breeding, or in pregnant or lactating bitches has not been evaluated. Please refer to the packaging insert for complete product information.

### **Meloxidyl® (meloxicam) Injection ISI for cats:**

**Repeated use of meloxicam in cats has been associated with acute renal failure and death. Do not administer a second dose of meloxicam. Do not follow meloxicam dose with any other NSAID.** As a class, cyclo-oxygenase inhibitory NSAIDs may be associated with gastrointestinal, kidney, or liver side effects. The most common side effects reported in field studies were vomiting and soft stool/diarrhea. These are usually mild, but may be serious. If side effects occur, pet owners should contact their veterinarian. Pets should be evaluated for pre-existing conditions and currently prescribed medications prior to treatment with MELOXIDYL, then monitored regularly while on treatment. Concurrent use with another NSAID, corticosteroid, or nephrotoxic medication should be avoided. The safe use of MELOXIDYL 5 mg/mL Solution for Injection in cats younger than 4 months of age, cats used for breeding, or in pregnant or lactating queens has not been evaluated. Please refer to the package insert for complete product information.

### **Meloxidyl® (meloxicam) Injection ISI combo (cat and dog):**

**Repeated use of meloxicam in cats has been associated with acute renal failure and death. Do not administer a second dose of meloxicam in cats. Do not follow meloxicam dose with any other NSAID in cats.** As a class, cyclo-oxygenase inhibitory NSAIDs may be associated with gastrointestinal, kidney, or liver side effects. The most common side effects reported in field studies were vomiting and soft stool/diarrhea. These are usually mild, but may be serious. If side effects occur, pet owners should contact their veterinarian. Pets should be evaluated for pre-existing conditions and currently prescribed medications prior to treatment with MELOXIDYL, then monitored regularly while on treatment. Concurrent use with another NSAID, corticosteroid, or nephrotoxic medication should be avoided. The safe use of MELOXIDYL in dogs younger than 6 months of age, dogs used for breeding, or in pregnant or lactating bitches has not been evaluated. The safe use of MELOXIDYL 5 mg/mL Solution for Injection in cats younger than 4 months of age, cats use for breeding, or in pregnant or lactating queens has not been evaluated. Please refer to the package insert for complete product information.