



October 1 - December 31, 2022

## Introducing the Newest Additions to Ceva's High-Quality Pain and Mobility Portfolio


**Meloxidyl®**  
(meloxicam)  
Injection 5 mg/mL

**Carprodyl®**  
(carprofen)  
50 mg/mL Injection

High quality products at a low cost!



- Non-steroidal anti-inflammatory drug for use in dogs and cats
- Controls osteoarthritis (OA) pain and inflammation in dogs
- Controls postoperative pain and inflammation in cats associated with orthopedic surgery, ovariectomy, and castration



- Non-steroidal anti-inflammatory drug for use in dogs only
- Controls relief of pain and inflammation associated with osteoarthritis (OA) in dogs
- Controls postoperative pain associated with soft tissue and orthopedic surgeries in dogs

Brand & Size	Product Number	Vet Price Per Unit	Special Introductory Price*
Meloxidyl® (meloxicam) Injection 10mL Vial	D65210I	\$66.98	<b>\$60.28</b>
Meloxidyl® (meloxicam) Injection 20mL Vial	D65220B	\$124.65	<b>\$112.18</b>
Carprodyl® (carprofen) Injection 20mL Vial	G43010B	\$64.20	<b>\$57.78</b>

\*Must purchase a minimum quantity of 2, mix or match, to qualify for the special introductory price. Purchase must be on a single invoice.

**Promotion terms:** Ceva reserves the right to vary the terms and conditions of this promotion or cancel this promotion without notice. Distributor has the right to make the final decision regarding all billing terms. To receive promotional offer, products must be purchased on the same order.

**Meloxidyl® (meloxicam) Injection 5 mg/mL - Repeated use of meloxicam in cats has been associated with acute renal failure and death. Do not administer a second dose of meloxicam in cats. Do not follow meloxicam with any other NSAID in cats.** As a class, NSAIDs may be associated with gastrointestinal, kidney, or liver side effects. The most common side effects reported in field studies were vomiting and soft stool/diarrhea. These are usually mild, but may be serious. If side effects occur, pet owners should contact their veterinarian. Pets should be evaluated for pre-existing conditions and currently prescribed medications prior to treatment with MELOXIDYL, then monitored regularly while on treatment. Concurrent use with another NSAID, corticosteroid, or nephrotoxic medication should be avoided. The safe use of MELOXIDYL in dogs younger than 6 months of age, dogs used for breeding, or in pregnant or lactating bitches has not been evaluated. The safe use of MELOXIDYL Injection in cats younger than 4 months of age, cats used for breeding, or pregnant or lactating queens has not been evaluated. Please refer to the package insert for complete product information.

**Carprodyl® (carprofen) Injection 50 mg/mL - For use in dogs only. Do not use in cats.** As a class, NSAIDs may be associated with gastrointestinal, kidney, or liver side effects. The most common side effects reported in field studies were vomiting and soft stool/diarrhea. These are usually mild but may be serious. If side effects occur, pet owners should contact their veterinarian. Pets should be evaluated for pre-existing conditions and currently prescribed medications prior to treatment with CARPRODYL, then monitored regularly while on treatment. Concurrent use with another NSAID or a corticosteroid should be avoided. The safe use of carprofen Injection in animals less than 6 weeks of age, pregnant dogs, dogs used for breeding purposes, or lactating bitches has not been established. Please refer to the package insert for complete product information.

